



FEBRUARY 2025

# Broccoli/cauliflower nutrition guide

## About this guide

The Soil Wealth ICP team has developed a series of crop nutrition guides to help growers, advisors and other vegetable industry service providers to manage crop-specific nutrition. The guides explore key considerations to ensure crop growth and nutrient needs are met, including:

- Growth stages of the crop and the nutritional focus for each stage
- Determining and calculating nutrient removal
- Factors affecting nutrient availability and uptake in the crop
- Approaches to crop monitoring, and
- Other helpful resources and information on nutrient functions.

Other nutrition guides include:

- Carrot: [soilwealth.com.au/2025/08/carrot-nutrition-guide/](http://soilwealth.com.au/2025/08/carrot-nutrition-guide/)
- Cucurbit: [soilwealth.com.au/2025/08/cucurbit-nutrition-guide/](http://soilwealth.com.au/2025/08/cucurbit-nutrition-guide/)
- Lettuce: [soilwealth.com.au/2025/08/lettuce-nutrition-guide/](http://soilwealth.com.au/2025/08/lettuce-nutrition-guide/)
- Onion: [soilwealth.com.au/2025/07/onion-nutrition-guide/](http://soilwealth.com.au/2025/07/onion-nutrition-guide/)

(developed by the Australian Onion Industry Communications and Extension Program (VN21000).

## CONTENTS

<b>About this guide</b> .....	<b>1</b>
<b>1. Broccoli and cauliflower growth and nutrient needs</b> .....	<b>2</b>
1.1 Broccoli and cauliflower growth stages and nutritional focus .....	4
<b>2. Crop monitoring</b> .....	<b>8</b>
2.1 Determining nutrient removal .....	10
2.2 Calculating nutrient removal .....	11
<b>3. Nutrition strategies and factors affecting nutrient availability and uptake for broccoli and cauliflower</b> .....	<b>13</b>
3.1 Factors affecting nutrient availability and uptake .....	13
<b>Other helpful resources</b> .....	<b>14</b>
Soil Wealth ICP Phase 2 outputs.....	14
Tools/calculators for nutritional needs ...	14
<b>References</b> .....	<b>14</b>
<b>Appendix</b> .....	<b>15</b>
Salinity case study .....	15
Nutrient functions .....	15

## 1. Broccoli and cauliflower growth and nutrient needs

When determining crop nutritional needs, it is important to be familiar with the site and local soil and climatic conditions as well as the cropping system. The following aspects should be considered:

- Crop, variety and growing days
- Growing season (weather impacts)
- Soil type and texture (approximate clay, silt, sand proportions)
- Soil condition/physical properties in the rootzone and beyond (e.g. compaction layers, other subsoil restrictions, salinity or sodicity, water holding capacity, drainage water and logging risks)
- Soil uniformity (across paddock, in the soil profile) – variable soil conditions may need targeted testing of management zones and variable application rates for best results
- Site history (crop rotations over the past 4-5 years, previous fertiliser applications, organic amendments such as composts, as well as pest, disease and weed history)
- Cultivation practices/tillage
- Irrigation system and practices
- Water quality
- Available and/or preferred nutrition products and soil amendments

- Application equipment
- Ability to adjust nutrition program during the season.

Having analytical data from comprehensive soil tests (including nutrient levels, CEC, pH, EC, organic carbon, total carbon and nitrogen) is a prerequisite to making decisions about applications of fertilisers or soil amendments. For more information refer to the Fertcare soil sampling guide and Soil Wealth ICP soil test interpretation guide (please refer to *Other helpful resources* on p.14).


The above-mentioned information and trends from soil tests taken over several previous years should be considered when making decisions about nutrient needs for a specific crop within the farming system.

- Soil tests are used to ensure that a base fertiliser application balances out deficiencies and imbalances in the soil as much as possible. For intensive crops like broccoli/cauliflower, annual soil testing is recommended.
- Knowing the expected yield and the crop growth stages (for topdressing and foliar applications) is critical to ensure the crop is getting the nutrients it needs at each phenological growth stage.





- Growth stages can be determined using the BBCH growth scale<sup>1</sup>. The following are 10 stages of plant growth in the BBCH scale<sup>2</sup>:
  1. Germination/early crop establishment
  2. Leaf development
  3. Formation of side shoots
  4. Stem elongation
  5. Development of harvestable vegetative plant parts
  6. Inflorescence emergence
  7. Flowering
  8. Development of fruit
  9. Ripening of fruit and seed
  10. Senescence.
- Nutrient removal rates based on the expected yield are used to ensure that the crop receives the amounts of nutrients it is expected to remove from the soil during its growth. The base fertiliser will provide some, or sometimes all the nutrients that the crop will remove for fast growing crops. For broccoli and cauliflower, a nutrient budget has to be developed that includes top dressing at key growth stages and monitoring approach (e.g. sap or dry tissue testing to ensure the program is working well).
- Nutrients that will not be easily lost and may be fixed by the soil (e.g. phosphorus) may be applied as a base dressing to cover the entire season. In phosphorus-fixing soils, top dressing may still be required. Phosphorus input to phosphorus-fixing soils have to be above the phosphorus removal rate. Local guidance from an experienced agronomist should be sought.
- Fast growing crops like baby leaf types, will usually not be top-dressed. They may receive a foliar application or fertigation if required to deal with imbalances.
- Slow growing crops like most head broccoli/cauliflower crops will require top-dressing at key growth stages. This is especially important for nutrients that can be easily lost from the rootzone such as nitrogen, sulphur, boron or potassium in sandy soils. Nutrients like potassium and nitrogen are needed in relatively large quantities and therefore spreading the nutrition program over the growing season to match nutrient uptake requirements at each growth stage is vital.




<sup>1</sup> [politicheagricole.it/flex/AppData/WebLive/Agrometeo/MIEPFY800/BBCHeng12001.pdf](http://politicheagricole.it/flex/AppData/WebLive/Agrometeo/MIEPFY800/BBCHeng12001.pdf)

<sup>2</sup> [eos.com/blog/stages-of-plant-growth/](http://eos.com/blog/stages-of-plant-growth/)


## 1.1 Broccoli and cauliflower growth stages and nutritional focus



Table 1 provides an overview of the nutritional aspects to consider for head broccoli/cauliflowers (e.g. cauliflower and broccoli) at key crop growth stages.

**Table 1:** Broccoli growth stages and nutritional focus





Crop stage	Image (photos or drawings)	Description	Comments	Focus
Pre plant		<p>Evaluate all site and local conditions and data mentioned in section 1 above this table.</p> <p>Determine total nutrient requirements based on soil tests and removal rates, whether the crop is grown from seed or transplanted to develop the crop nutrient budget.</p> <p>If an adviser, consider the operational aspects of the site in discussion with the grower.</p>	<p>The main purpose of preplant fertiliser is to ensure the soil has no nutrient deficiencies or imbalances.</p> <p>It is important to consider crop and water relations (soil texture/conditions and irrigation practices) for nutrient management of the rootzone.</p> <p>Ensure that the soil pH levels are optimal for nutrient uptake and to ensure that the environment is not conducive to disease.</p>	<p>Site conditions and suitability for crop.</p> <p>Prepare the soil to ensure the physical properties are adequate for nutrient availability with good rooting depth and no compaction.</p> <p>Ensure that fertiliser is applied for optimal nutrient use efficiency. For example, phosphorus should be applied as a broadcast or banded as close to planting as possible in phosphorus-fixing soils.</p> <p>Using nitrogen fertilisers with denitrification inhibitors is one way of reducing nitrogen top dressing needs.</p> <p>Brassica crops require good soil levels of sulphur and boron.</p>



Crop stage	Image (photos or drawings)	Description	Comments	Focus
<p>At planting and through to germination (direct seeded crop), early crop establishment (BBCH stage 1)</p>		<p>The nutrient application should be designed to last until the first planned topdressing.</p> <p>Ensure there is a small surplus amount of nutrients to ensure that the crop is not deficient in nutrients over the course of the growth cycle. The soil pH has to be above pH 7 to reduce the risk of clubroot infections.</p>	<p>Saline (high EC/chloride) and sodic soils may require higher levels of calcium and potassium than soils without these problems. The same applies if irrigation water is saline.</p> <p>Crops grown in poorly structured soil will have a restricted root zone. Therefore, they may require feeding more frequently with smaller amount of nutrients than crops grown in well-structured soil.</p> <p>Sandy soils and those with low organic matter levels may also need fertilising more often with smaller amounts of nutrients than soil with good organic matter levels and/or heavier soils.</p>	<p>Compound fertilisers as a banded or broadcast application are a good option to ensure even distribution of essential elements, especially calcium and trace elements.</p> <p>Mixed fertilisers easily separate in the spreader and especially trace element application will be uneven.</p> <p>Foliar applications may be a good option, if compound fertilisers are not used.</p>

Crop stage	Image (photos or drawings)	Description	Comments	Focus
<p>Leaf development and vegetative growth (BBCH stages 2-5)</p>		<p>Plant nutrient tests should be performed from early growth stages and throughout vegetative growth to allow time for nutrient imbalances to be corrected.</p> <p>Once head development is well underway, testing may be used for diagnostic purposes, but correcting nutrient deficiencies will become more difficult as the plant ages.</p>	<p>The youngest fully expanded leaf should be sampled for dry matter testing.</p> <p>For sap testing, the leaf petioles of these leaves are good indicators of the plant's most recent nutrient uptake.</p> <p>If planning for 1-2 tests only, focus on the earlier stages when imbalances are easier to correct.</p>	<p>Nitrogen is important for early development when plants are growing rapidly, but must be in balance with other nutrients, especially calcium.</p> <p>Calcium is critical for early head development, strong structure, and shelf life. It is best applied in good time to be available when the head starts to form. Foliar applications are possible.</p>
<p>Head development (BBCH stage 6) Early to mid head development</p>		<p>If calcium is not applied prior to this stage, foliar applications should be considered.</p> <p>Plant sap analysis can still be used at early head formation to amend any nutrient requirements.</p>	<p>Calcium nitrate and boron can be used to help maintain a good rootzone environment for optimal nutrient uptake and to combat adverse saline (high soil sodium chloride) or sodic (high sodium) conditions.</p>	<p>Nitrogen should best be applied in the plant available form as nitrate if immediate plant uptake is important.</p> <p>Trace elements are now best applied with foliar applications. Foliar fertilisers may also help to correct low plant levels of potassium and calcium.</p>



Crop stage	Image (photos or drawings)	Description	Comments	Focus
<p>Head development (BBCH stage 7-8)</p> <p>Mid-head development to harvest</p>	 	<p>Potassium demand is at its highest during head development.</p> <p>In most plants, potassium is removed at a faster rate than nitrogen.</p> <p>Monitor nutrient levels with sap tests if required for diagnostic purposes.</p>	<p>Potassium can be applied as a liquid thiosulphate or as potassium sulphate or potassium nitrate, if nitrogen is needed.</p> <p>Potassium may be limiting in soils, even if it is at satisfactory levels because its rate of release from the soil cannot keep up with the demand of the crop during a rapid phase of growth.</p>	<p>Apply potassium before the period of rapid growth (at mid head development).</p> <p>Ensure adequate uptake of calcium and potassium while maintaining magnesium and trace elements.</p>
<p>Harvest (BBCH 9)</p>	 	<p>It is too late to make any nutrient amendments.</p>	<p>To ensure the site is restored for the next crop, consider soil nutrient testing and the nutrients that will be returned to the soil with crop residues to ensure that nutrient levels are replenished to the desirable crop requirement but not oversupplied.</p> <p>For instance, residues from a broccoli or cauliflower crop can return &gt;170 kg/nitrogen per hectare to the soil.</p>	<p>Plan soil nutrient testing ahead of the next crop for the site.</p>



## SOME KEY CONSIDERATIONS

- Experience and knowledge of soil conditions and weather
- Yield potential and crop removal figures
- Growth stages and expected rooting depth and distribution
- Nutrient uptake efficiency and limiting factors
- Soil management, mulching and green crops may have a greater impact on nutrient uptake than fertiliser applications
- Growing conditions and irrigation management may have a greater impact on nutrient uptake than fertiliser applications
- Stress, including from pests and diseases, will influence nutrient uptake significantly
- Good decisions are based on good data such as soil, plant and water test results.

## 2. Crop monitoring

**While a soil analysis provides information about soil reserves, deficiencies and imbalances in the soil and potential nutrient availability, a plant analysis provides information about actual nutrient uptake and use.**

Plant analyses are required to gain information about the nutritional status of a crop before problems are visible. If symptoms appear, plant analysis can be used to diagnose nutrition problems (deficiencies, imbalances, toxicity) as early as possible. Planned, repeated monitoring within a crop cycle helps to assess nutrient availability within the soil and removal of key nutrients by a crop. Plant analysis data can be used to replace nutrients in the soil after harvest.

### Procedures for plant analysis include:

- Plant sap testing – a quick way to check the nutritional status of your crop. Sap testing offers early detection and correction of nutritional problems in the crop.
- Plant dry matter (or tissue) analysis – as a measure of nutrient accumulation in the tissue, the analysis can provide information for checking the effectiveness of nutrient budgeting, calculation of crop nutrient removal (if the harvested plant part is analysed) or total nutrient uptake of biomass samples.

## WHICH ANALYSIS TO CHOOSE

- Determining which test to do is based on how the data is to be used
- For nutrient budgeting based on removal post-harvest, a dry matter (tissue) test of the harvested plant parts is required
- For monitoring nutrient uptake during the growth cycle, a plant sap test is more suitable
- To correct nutrient deficiencies, samples should be taken during early growth, well before symptoms appear.

If nutrition trends over time are monitored, it is recommended to take a sap test at a consistent time of day when transpiration rates are low and comparable. Never sample stressed plants for any type of analysis.

### Sampling

- Plant dry matter
  - Sample the entire youngest fully expanded leaf (monitoring)
  - Sample the entire heads at harvest (nutrient removal)
- Plant sap
  - Sample the petiole in the youngest fully expanded leaf.

## WHAT TO CONSIDER FOR A FERTILISER PROGRAM

- Nutrient removal rates at the expected yield level
- Crop requirements at different growth stages
- Rooting depths, root distribution
- Crop and variety specific needs
- Expected weather conditions
- Water quality and irrigation system
- Available/preferred application methods and fertilisers
- Monitoring plan – actual nutrient uptake during early growth
- Ability to adjust nutrition program during the season.

For more information on crop monitoring please refer to the Soil Wealth ICP Plant analysis for vegetable crops guide:

[soilwealth.com.au/2020/02/plant-analysis-for-vegetable-crops-a-practical-guide-to-sampling-analysis-and-interpretation/](https://soilwealth.com.au/2020/02/plant-analysis-for-vegetable-crops-a-practical-guide-to-sampling-analysis-and-interpretation/)

For sampling procedures please refer to the Fertcare Soil Sampling Guide:

[fertilizer.org.au/Portals/0/Documents/Fertcare/Fertcare%20Soil%20Sampling%20Guide.pdf?ver=2019-06-17-095413-863](https://fertilizer.org.au/Portals/0/Documents/Fertcare/Fertcare%20Soil%20Sampling%20Guide.pdf?ver=2019-06-17-095413-863)

For information on soil test interpretation please refer to the Soil Wealth ICP Soil testing and interpretation for vegetable crops guide:

[soilwealth.com.au/2017/05/soil-testing-and-interpretation-for-vegetable-crops-a-guide/](https://soilwealth.com.au/2017/05/soil-testing-and-interpretation-for-vegetable-crops-a-guide/)

## WHEN INTERPRETING SOIL TEST RESULTS, CONSIDER:

- Mobility/movement of different nutrients, nutrient interactions e.g. cations, phosphorus and zinc, trace metals, and electrical conductivity (EC) and pH
- Specific nutrient needs of the crop
- Soil-related constraints such as salinity or sodicity, rootzone restrictions (e.g. compaction, sudden changes in texture) that limit root growth.



## 2.1 Determining nutrient removal

During harvest, nutrients are removed from the system. If the soil is in balance, only what is removed with harvest needs to be replaced via mineral or organic fertilisers. Thus, crop removal figures can be used as a starting point for nutrient budgeting. Various nutrient budgeting tools are available (see Resources on p.14).

1. Use a soil test to determine nutrient deficiencies and imbalances in the soil (best over the main rootzone depth). Calculate the amount of nutrients required to balance the soil, so that all nutrients are within a satisfactory range in the soil.
2. To convert mg/kg to kg/ha, multiply the soil test value in each sampling layer by the depth of the soil sample (millimetres), then multiply by the bulk density (g/cm<sup>3</sup>) and divide by 100, then add up the values for each layer to arrive at nutrient content in kg/ha to the total soil depth sampled.

Explanation and example for converting mg/kg (=ppm) of a nutrient in a soil test to kg/ha of that nutrient for a sampled soil depth:

**Calculation: kg/ha of selected nutrient = Soil test value (mg/kg) × Bulk density (g/cm) × Soil depth (m) × 10.**

### Step-by-step explanation


- mg/kg (ppm) from soil test – this is the concentration of the nutrient in the soil.
- Bulk density (g/cm<sup>3</sup>) – typical values:
  - Sandy soil: ~1.4 g/cm<sup>3</sup>
  - Loamy soil: ~1.3 g/cm<sup>3</sup>
  - Clay soil: ~1.2 g/cm<sup>3</sup>
  - Soil depth (cm) – e.g., 20 cm (= 0.2 m).
  - Factor (10) – to convert g/m<sup>2</sup> to kg/ha.

Example calculation

<b>Soil test value</b>	<b>50</b>	<b>mg/kg</b>
<b>Bulk density</b>	<b>1.3</b>	<b>g/cm<sup>3</sup></b>
<b>Sampling depth</b>	<b>0.2</b>	<b>m</b>
<b>Factor</b>	<b>10</b>	
<b>Result</b>	<b>130</b>	<b>kg/ha</b>

If a soil test shows 50 mg/kg (ppm) of a nutrient in a loamy soil (bulk density = 1.3 g/cm<sup>3</sup>) sampled to 0.2 m depth:

kg/ha = 50 × 1.3 × 0.2 × 10 = 130 kg/ha of nutrient to 20 cm soil depth.



## 2.2 Calculating nutrient removal



Nutrient removal is the amount of all nutrients removed at harvest (total yield taken off without trimming) in kilograms per tonne (kg/t) of dry matter removed.

For a nutrient budget, the anticipated total yield in tonnes per hectare (t/ha) is used to estimate nutrient removal by the crop. This helps to determine how much of each nutrient is required to grow the crop (kg/ha), if the soil is in balance (i.e. has no deficiencies). The principle is to provide the nutrients that will be removed over the growing season.

Example of calculating nitrogen removal rates:

<b>60,000</b>	<b>heads/ha planted</b>
<b>0.6</b>	<b>kg/head harvested, not trimmed</b>
<b>36,000</b>	<b>kg total removed from the paddock</b>
<b>10</b>	<b>% dry matter (DM)</b>
<b>3,600</b>	<b>kg DM/ha removed</b>
<b>4.50%</b>	<b>nitrogen in head dry matter (based on a dry tissue analysis)</b>
<b>162.00</b>	<b>kg nitrogen removed with harvest</b>

Figure 1 provides an example of nutrient removal rates for broccoli and cauliflower in kg/tonne. The examples show that broccoli removes more nitrogen than cauliflower. The nutrients in the crop residue will be returned to the soil and become available to the following crop or cover crop as the biomass breaks down. This should be considered in the nutrient budget for the following crops.



**Figure 1:** Example of nutrient removal rates for cauliflower and broccoli. Major elements are shown in kg/tonne harvested crop. (N: nitrogen, P: phosphorus, K: potassium, S: sulphur, Ca: calcium, Mg: magnesium.)

**Replenishment of nutrients based on removal rates**

<b>N</b>	Nitrogen: Match removal rates, reduce for high organic matter soils, potentially increase for depleted soils, or very light soils	3–4 split applications according to growth stages and plant testing results, broccoli needs little nitrogen early in its growth; excess nitrogen applied early is likely to be lost from the rootzone
<b>P</b>	Phosphorus: 2–3 x removal rate, potentially more on phosphorus-fixing soils	Phosphorus can be pre-spread; phosphorus top dressing or foliar applications may be required on phosphorus-fixing soils based on plant testing results
<b>K</b>	Potassium: Match removal	Split applications are recommended for light textured soils, topdressing should occur before major growth phases (high biomass production, head development); potassium is commonly required at similar rates as nitrogen
<b>Ca</b>	Calcium: Up to 2 x removal rate, reduce in high calcium soils	
<b>Mg</b>	Magnesium: Match removal, reduce in high magnesium soils	
<b>S</b>	Sulphur: Match removal, increase for depleted soils	Consider sulphur proportion in magnesium and potassium fertilisers

The above approach is based on adding nutrients that correspond with removal rates to a balanced soil with nutrients within the desirable range or 'topped up' to the desirable range as part of a pre plant application.

**Calculating fertiliser application rates**

kg/ha nutrient required ÷ % nutrient in fertiliser x 100 = amount of fertiliser kg/ha to apply.



### 3. Nutrition strategies and factors affecting nutrient availability and uptake for broccoli and cauliflower

Refer to the Soil Wealth ICP nutrition products guide and the soil test interpretation guide (see Resources on p.14) for more information about nutrients in fertilisers and soil amendment as well as nutrient interactions and their movement in soils and plants. Crop nutrition should consider the following factors that affect the availability of nutrients to plants.

#### 3.1 Factors affecting nutrient availability and uptake

- **Physiological disorders** – many nutritional disorders are initiated as early as during seed germination. Therefore, it is important to have a relationship with the supplier of seedlings to ensure they are of the required quality and nutrient level/balance.
- **Soil salinity and sodicity** influence the quantity of cations (potassium, magnesium, calcium) that can be taken up by the plant due to cation competition with sodium. High sodium levels in the soil have a negative effect on soil structure and therefore affect root health and nutrient movement.
- **Soil structure** – compaction leads to poor root penetration, water infiltration and drainage which reduces nutrient uptake.

- **Organic matter** affects nutrient availability and uptake positively as it increases porosity (air spaces and water infiltration, storage and drainage); it provides food for soil biology, a high cation exchange capacity and source of nutrients via mineralisation.
- **Soil biology/microbial composition** is important for improving nutrient cycling and plant nutrient uptake.
- **Soil acidity or alkalinity (pH)** – soils can become more acidic or alkaline depending on the type of fertiliser used, which affects nutrient uptake. Nitrogen is the main nutrient affecting pH. Ammonium-based products and urea have the greatest potential to acidify soil. Lime or dolomite can be applied to correct soil pH; dolomite should be used when magnesium is low and lime if calcium is low in the soil. Gypsum can be applied to supply calcium and sulphur to the soil but it does not affect the soil pH.



## Other helpful resources

### Soil Wealth ICP Phase 2 outputs

1. Getting the soil pH right – Lime quality and application rates (fact sheet, January 2019)  
[soilwealth.com.au/2019/01/getting-soil-ph-right-lime-quality-and-application-rates/](http://soilwealth.com.au/2019/01/getting-soil-ph-right-lime-quality-and-application-rates/)
2. Managing salinity in vegetable crops (fact sheet, August 2019)  
[soilwealth.com.au/2019/08/managing-salinity-in-vegetable-crops/](http://soilwealth.com.au/2019/08/managing-salinity-in-vegetable-crops/)
3. Soil phosphorus – The basics (fact sheet, September 2019)  
[soilwealth.com.au/2019/09/soil-phosphorus-the-basics/](http://soilwealth.com.au/2019/09/soil-phosphorus-the-basics/)
4. Nitrate field test (fact sheet, January 2020)  
[soilwealth.com.au/2020/01/nitrate-field-test/](http://soilwealth.com.au/2020/01/nitrate-field-test/)
5. Taking soil samples (fact sheet, January 2020)  
[soilwealth.com.au/2020/01/taking-soil-samples/](http://soilwealth.com.au/2020/01/taking-soil-samples/)
6. Plant analysis for vegetable crops: A practical guide to sampling, analysis and interpretation by Doris Blaesing, RMCG (Version 2, February 2020)  
[soilwealth.com.au/2020/02/plant-analysis-for-vegetable-crops-a-practical-guide-to-sampling-analysis-and-interpretation](http://soilwealth.com.au/2020/02/plant-analysis-for-vegetable-crops-a-practical-guide-to-sampling-analysis-and-interpretation)
7. Managing sodicity in vegetable crops (fact sheet, May 2020)  
[soilwealth.com.au/2020/05/managing-sodicity-in-vegetable-crops/](http://soilwealth.com.au/2020/05/managing-sodicity-in-vegetable-crops/)
8. Soil phosphorus – The basics (fact sheet, September 2019)  
[soilwealth.com.au/2019/09/soil-phosphorus-the-basics/](http://soilwealth.com.au/2019/09/soil-phosphorus-the-basics/)
9. Nutrition management resources (fact sheet, June 2020)  
[soilwealth.com.au/2020/06/nutrition-management-resources/](http://soilwealth.com.au/2020/06/nutrition-management-resources/)
10. Soil testing and interpretation for vegetable crops by Doris Blaesing, RMCG (May 2017)  
[soilwealth.com.au/2017/05/soil-testing-and-interpretation-for-vegetable-crops-a-guide/](http://soilwealth.com.au/2017/05/soil-testing-and-interpretation-for-vegetable-crops-a-guide/)
11. Crop nutrition products – A guide to product types, properties and uses  
*Coming soon*

### Tools/calculators for nutritional needs

1. Nutrient removal calculator  
[ipni.net/article/IPNI-3346](http://ipni.net/article/IPNI-3346) or [fertilizer.org](http://fertilizer.org)
2. Fertcare Soil Sampling Guide  
[fertilizer.org.au/Portals/0/Documents/Fertcare/Fertcare%20Soil%20Sampling%20Guide.pdf?ver=2019-06-17-095413-863](http://fertilizer.org.au/Portals/0/Documents/Fertcare/Fertcare%20Soil%20Sampling%20Guide.pdf?ver=2019-06-17-095413-863)

### References

1. International Fertiliser Association (IFA)  
[fertilizer.org](http://fertilizer.org)
2. AgVita Analytical  
[agvita.com.au](http://agvita.com.au)
3. Campbells Fertilisers Australia  
[campbellsfert.com.au](http://campbellsfert.com.au)



## Appendix

### Salinity case study

Many Australian soils are affected by salinity. Werribee South located 32 km south-west of Melbourne is a large vegetable growing region covering an area of 3,000 hectares which consists of a high proportion of lettuce and broccoli/cauliflower production. The Werribee South growing region is part of the Werribee Irrigation District Recycled Water Scheme which provides irrigation water to a large proportion of vegetable farms. The irrigation water is relatively high in salt concentration, containing particularly high levels of sodium and chloride. This high salt level in the water results in high electrical conductivity which fluctuates between 800 mg/L and more than 1000 mg/L, an equivalent EV of 1.4 dS/m to 1.8dS/m. The low rainfall of 542 mm means that the salt is not washed away from the rootzone.

A demonstration trial was conducted at Fresh Select, a vegetable business in Werribee South, to examine the effects of different soil ameliorants including compost, gypsum and calcium thiosulphate on soil and plant sodium levels. The plant sap tests showed that the plants that received the calcium thiosulphate treatment had lower sodium levels in combination with compost compared to the control bay which had

only received gypsum. Calcium thiosulphate displaces the sodium from the soil exchange sites. The thiosulphate ion slowly oxidises releasing additional calcium for greater sodium removal compared with gypsum which contains sulphur and has low solubility and an inconsistent release of calcium.

### Nutrient functions

#### Macro nutrients (major elements)

##### **Nitrogen**

Nitrogen (N) is an essential component of amino acids for building proteins, nucleic acids and chlorophyll for photosynthesis. Nitrogen is available to the plant in the form of nitrate which in combination with ammonia should not exceed 50 kg/ha in the root zone. Nitrogen use efficiency (NUE: % = N removed/ N applied x 100) is calculated after harvest and shows how much N has been used by the crop.

The key growth stages for nitrogen requirement are during intensive leaf production. Nitrogen requirements will vary between broccoli/cauliflower types and range between 1.6kg/t – 4.7 kg/t.

##### **Phosphorus**

Readily available phosphorus (P) is important for early growth stages to promote root development and to boost establishment in broccoli/cauliflower crops through

the vegetative growth stages. Australian soils are low in P which is non-labile (not available for plant uptake). Given the nature of phosphorus availability from soils, P replacement rates need to be higher (2-3 times) more than removal rates especially in heavier P-fixing soils that are high in aluminium and iron (low pH) and high in calcium (high pH). The most common phosphorus sources are superphosphate-based fertilisers (e.g. single superphosphate, SSP 7-9% P or triple superphosphate, TSP 44-46% P). The key difference between single superphosphate and triple superphosphate is that single superphosphate is produced from phosphate rock and sulfuric acid, whereas triple superphosphate is produced from phosphate rock and phosphoric acid. Humic acids, monopotassium phosphate (MKP) or liquid P fertiliser formulations can help acidify the rootzone to make P more readily available. Nitrogen and phosphorus can provide a synergistic effect when banded together, N providing an increased amount of P uptake. Good soil structure and organic matter will support microbial activity and mycorrhiza association with the roots which will improve P availability and uptake. Broccoli/cauliflower utilise around 0.33 -0.67 kg P/t of total fresh matter.

##### **Potassium**

Potassium (K) is required for plant water relations by regulating stomatal functions and water use efficiency in plant transpiration. In broccoli/cauliflower crops potassium is in high demand during the mid to late heading stages. Potassium provides strength to plant cell walls enhancing crop vigour, helps form and remove starches, sugars and oils and decreases susceptibility to diseases and insects. K can be applied in the forms of muriate of potash (potassium chloride) and sulphate of potash (potassium sulphate). The common sources of K fertilisers include potassium nitrate and potassium thiosulphate, and mono ammonium phosphate (MAP). Depending on the broccoli/cauliflower type, between 2.7-5.1 kg/t of K is required with significant proportions removed in cabbage crops.

##### **Calcium**

Calcium (Ca) is essential for root health, growth of new roots and root hairs, and the development of leaves as well as building strong cell walls. Adequate calcium levels in the plant are important for good storage performance.

Calcium uptake is driven by transpiration, i.e., the loss of water through leaves (passive transport). This means that calcium only



moves upwards with water, not from leaves to roots (no active transport). Calcium also protects the plant against diseases (fungi and bacteria), regulates stomata aiding with transpiration and contributes to crop quality. The soil pH influences calcium availability, generally more alkaline soil increases the availability of calcium, hence calcareous soils in Victoria's Mallee region. Calcium competes with other cations (positively charged ions). Saline soils can impact the uptake of calcium as sodium (Na<sup>+</sup>) displaces calcium.

Ca is generally in short supply in acid soils. Lime, gypsum, dolomite and superphosphate (a mixture of calcium phosphate and calcium sulphate) all supply calcium to the soil but do not provide readily available Ca to the soil solution. Lime and dolomite raise the pH, gypsum improves soil structure in clay soils but does not change the pH. Peak calcium requirements is at the beginning of head formation. Most calcium is utilised for leaf growth with proportionally lower levels found in the harvested heads. Higher calcium tissue levels improve crop quality and storage.

Lime is often seen as the cheapest option for increasing soil Ca levels, and it is important for increasing the pH of acid soils (via its carbonate component), improving soil structure and microbial activity (as part of the pH and structure effect). However, soluble Ca should be provided especially

early in the season, e.g. via liquid fertilisers or calcium nitrate. When using calcium nitrate, care must be taken to not oversupply nitrate N to young crops when uptake is low. Calcium nitrate may be a good topdressing fertiliser.

### **Magnesium**

Magnesium (Mg) is a key component of the green pigment chlorophyll. It is vital for photosynthesis (the conversion of the sun's energy to energy for the plant). Magnesium is required at lower levels in broccoli/cauliflower than in other crops and removals are between 0.1-0.3kg/t depending on the broccoli/cauliflower crop. Deficiencies occur mainly on sandy, acid soils in high rainfall areas, especially if used for intensive horticulture or dairying. Heavy applications of potassium in fertilisers can also produce magnesium deficiency due to cation competition.

Magnesium deficiency in soils can be overcome with dolomite (magnesium-calcium carbonate, slow availability), magnesite (magnesium oxide) or Epsom salts (magnesium sulphate).

### **Micronutrients (trace elements)**

#### **Boron**

Boron (B), together with calcium, helps with the formation of cell walls in rapidly growing tissue. Deficiency reduces the uptake of calcium and inhibits the plant's ability

to use it. B is one of the most prevalent micronutrient deficiencies in Australian soils used for horticulture.

B can be reasonably easily applied to the soil via borax or calcium nitrate plus B. Do not exceed the recommended amount since B can be toxic. Applications need to be based on soil testing results. Foliar applications work as well, and rates must not exceed label recommendations because B will be phytotoxic if oversupplied. Plant testing can provide information on how much to apply.

B uptake is driven by transpiration, i.e., the loss of water through leaves (passive transport). This means that B only moves upwards with water, not from leaves to roots (no active transport).

However, soil plant testing should be used to confirm the need for B fertilisers to avoid phytotoxicity.

#### **Zinc**

Zinc (Zn) helps in the production of a plant growth regulators responsible for stem elongation and leaf expansion. It is readily available in acid soils but is easily outcompeted for uptake by iron in volcanic soils. If Zn levels determined by soil testing are low addition of zinc sulphate can be used to increase soil levels. Zn is usually added in the preplant fertiliser. Crops can be sprayed with foliar Zn fertilisers if required based on plant testing. Chelated products will be most

efficient.

Excessive amounts of Zn can be toxic, so apply only if needed. Zn can compete with other trace metals (e.g. copper, manganese) for uptake.

#### **Copper**

Copper (Cu) is an essential constituent of enzymes in plants and can be readily available in some soils, although it can be deficient, especially in red soils. Cu can be applied as copper chelate EDTA, copper sulphate pentahydrate or copper sulphate monohydrate. Overuse of another trace element, e.g. molybdenum, can cause Cu deficiency in animals. Overuse can cause toxicity and reduce uptake of other trace metals (e.g. Mn).

#### **Iron**

Iron (Fe) is a constituent of many plant compounds that regulate and promote growth. It usually is readily available in naturally acid soils. Deficiencies can be addressed via foliar applications. These are commonly applied as a complete trace element mix.

#### **Manganese**

Manganese (Mn) helps with photosynthesis. It is freely available in naturally acid soils, often in toxic amounts in very acid soils, but can be deficient in sandy soils. Toxicity is remedied with lime.



### ***Molybdenum***

Molybdenum (Mo) helps bacteria and soil organisms convert nitrogen from the air to soluble nitrogen compounds in the soil, so is particularly needed by legumes. It is also essential in the formation of proteins from soluble nitrogen compounds taken up from the soil or leaves.

Mo deficiency is prevalent in many acid soils but can be remedied easily with applications of Mo super, molybdenum trioxide (applied during inoculation and lime pelleting of legume seed), or sodium molybdate (sprayed on emerging and young plants).

*For analysis of nutrients in soil testing refer to the Soil testing and interpretation for vegetable crops guide published as part of the Soil Wealth ICP project (see Resources on p.14).*



Hort Innovation, Applied Horticultural Research Pty Ltd (AHR) and RM Consulting Group (RMCG) make no representations and expressly disclaims all warranties (to the extent permitted by law) about the accuracy, completeness, or currency of information in this fact sheet. Users of this material should take independent action before relying on its accuracy in any way. Reliance on any information provided by Hort Innovation, AHR or RMCG is entirely at your own risk. Hort Innovation, AHR or RMCG are not responsible for, and will not be liable for, any loss, damage, claim, expense, cost (including legal costs) or other liability arising in any way (including from Hort Innovation, AHR, RMCG or any other person's negligence or otherwise) from your use or non-use of information from project MT22004 - Soil Wealth and Integrated Crop Protection - Phase 3 or from reliance on information contained in this material or that Hort Innovation, AHR or RMCG provides to you by any other means.